

November 8, 2022

Webinar ID  
819 4240 9471

#	Question	Answer
1	Getting my question in obnoxiously early-I've seen some studies lately saying vaginal estrogen isn't safe for patients on aromatase inhibitors. Any thoughts or data on this?	Vaginal estrogen tablets and ring are safe and effective, we don't recommend creams, and data is still emerging on prasterone.
2	Are there still the same contraindications when prescribing estrogen therapy to menopausal patients are there are when prescribing estrogen therapy to non-menopausal patients? (Example: smoking, migraines with aura, HTN,	No there is no since its a different estrogen. If they have known CVD we don't recommend estrogen.
3	I want to learn the dose and duration of local estrogen and testosterone for hormone induced vestibulydina	Differnt doses are used, estradiol/testosterone 0.01/0.1% to 0.03/0.1% are typical.
4	To clarify, is concomitant progesterone not indicated with local estradiol?	Yes, not needed.
5	Dr. Goldstein, the before and after pics are after local DHEA? Is that what you are using for your androgens here?	
6	Source for vaginal DHEA? compounded? Dose?	Intrarosa, FDA approved 6.5 mg daily insert.
7	Any tips for Intrarosa compliance? My patients hate that it's qd. Any new DHEA products on the pipeline?	
8	Do you routinely measure hormonal levels? Do you adjust therapy to number? Symptoms? Physical exam?	
9	Would it be beneficial to add DHEA to estradiol.01% and Testosterone.1%cream used for hormonally associated vestibulodynia and/or GSM? If so what dose?	
10	I have two question:	
11	1. Primarin cream is not available in our country. So can i use oestrogen (estradiol)? Is there any difference in the effectiveness of the treatment?	Similarly effective
12	2. When treating hypoactive sexual desire disorder, do you often use flibansetin or bremelanotide? If not, what can be replaced with this medicine?	This seminar was on GSM, not HSDD.